

Hausdorff dimension of the set of extreme points of a self-similar set

Andrew Tetenov and Ivan Davydkin ,

(*Gorno-Altai state university*)

Abstract

If the system of contracting similitudes on $\mathbb{R}^2$ satisfies open convex set condition, then the set of extreme points of the convex hull $\tilde{K}$ of it's invariant self-similar set has Hausdorff dimension 0 . If, additionally, all the rotation angles α_i of the similitudes φ_i are commensurable with π , then the set $\tilde{K}$ is a convex polygon.

1 Introduction

Let $\varphi_1, \dots, \varphi_n$ — be contracting similitudes on $\mathbb{R}^2$ with rotation angles α_i . Let $K = \bigcup_{i=1}^n \varphi_i(K)$ — be compact self-similar set with respect to the system $\{\varphi_1, \dots, \varphi_n\}$.

Let $\tilde{K}$ — be the convex hull $H(K)$ of the set K .

The boundary $\partial\tilde{K}$ of the set $\tilde{K}$ may be viewed as consisting of two parts: a compact set F of all extreme points of $\tilde{K}$, which we call *vertices* of $\tilde{K}$, and of an union of finite or countable family of open line intervals l_i , called *the sides* of $\tilde{K}$.

There is a number of well-known cases when the set $\tilde{K}$ is a convex finite-sided polygon. In Example 1 we show how mappings φ_i with irrational rotation angles α_i may produce self-similar sets K , whose convex hull $\tilde{K}$ has infinite set of sides (and hence of vertices) . So the question arises , what conditions must be imposed upon $\{\varphi_1, \dots, \varphi_n\}$ to obtain the set $\tilde{K}$ with finite number of sides and vertices and what is the structure of $\partial\tilde{K}$ in the case when the set F or $\{l_i\}$ is infinite.

To give a partial answer to these questions, we formulate open convex set condition (Definition 2), implying some finiteness properties for $\partial\tilde{K}$. Then we show that, if the system $(\varphi_1, \dots, \varphi_n)$ satisfies open convex set condition, then the set of vertices of $\tilde{K}$ has Hausdorff dimension 0 (Theorem 9). If, additionally, all the rotation angles α_i of the maps φ_i are commensurable with π , then the set of vertices of $\tilde{K}$ is finite, so the set $\tilde{K}$ is a convex polygon (Corollary 15).

The authors express their appreciation to V.V.Aseev who suggested the problem.

2 Preliminaries.

Let S be a system of contraction similitudes $\varphi_1, \dots, \varphi_n$, (written in the complex form as $\varphi_i(z) = q_i(z - z_i)e^{i\alpha_i} + z_i$, $0 < q_i \leq q < 1$, $\alpha_i \in [0, 2\pi]$). The system S defines the Hutchinson transformation $T(A) = \bigcup \varphi_i(A)$ on the space $C(\mathbb{R}^2)$ of compact non-empty subsets of $\mathbb{R}^2$. The transformation T is a contraction map on the space $C(\mathbb{R}^2)$ in the Hausdorff metrics. Compact invariant set $K = \bigcup_{i=1}^n \varphi_i(K)$ is a fixed point of the transformation T .

By $H(X)$ we denote the convex hull of a set X .

We shall denote by $\tilde{K}$ the convex hull $H(K)$ of the invariant set K with respect to the system S .

Using notation similar to that of [2], we denote

$$\varphi_{i_1 \dots i_m} = \varphi_{i_1} \circ \varphi_{i_2} \circ \dots \circ \varphi_{i_m}(K),$$

$$K_{i_1 \dots i_m} = \varphi_{i_1 \dots i_m}(K), \quad \tilde{K}_{i_1 \dots i_m} = \varphi_{i_1 \dots i_m}(\tilde{K}).$$

By $Q_{i_1 \dots i_m}$ we denote $\tilde{K}_{i_1 \dots i_m} \cap \partial \tilde{K}$ and

by $F_{i_1 \dots i_m} = \tilde{K}_{i_1 \dots i_m} \cap F^-$ the set of extreme points belonging to $\tilde{K}_{i_1 \dots i_m}$.

Let $\tilde{C}(\mathbb{R}^2)$ be the set of all convex compact subsets in $\mathbb{R}^2$. Define the transformation $\tilde{T}$ on the space $\tilde{C}(\mathbb{R}^2)$ by: $\tilde{T}(A) = H(T(A)) = H(\bigcup \varphi_i(A))$ for any $A \subset \tilde{C}(\mathbb{R}^2)$.

PROPOSITION 1 . *Let $S = \{\varphi_1, \dots, \varphi_n\}$ — be a set of contracting similitudes on $\mathbb{R}^2$ with $\text{Lip}(f_i) = q_i$, $0 < q_i \leq q < 1$. The transformation $\tilde{T}$ is a contraction map of the space $\tilde{C}(\mathbb{R}^2)$ in Hausdorff metrics, and the set $\tilde{K}$ is it's fixed point.*

Proof. For any two compact sets U_1, U_2 the Hausdorff distance between their convex hulls $d(H(U_1), H(U_2))$ is less or equal then $d(U_1, U_2)$. Since $\tilde{T}(A) = H(T(A))$, we obtain that $d(\tilde{T}(A), \tilde{T}(B)) \leq d(T(A), T(B)) < qd(A, B)$.

The equality $\tilde{T}(\tilde{K}) = H(\bigcup \varphi_i(\tilde{K})) = H(\bigcup \varphi_i(K)) = \tilde{K}$, shows that $\tilde{T}$ leaves $\tilde{K}$ fixed.

Example 1.

Consider a system $\varphi_1(z) = \frac{1}{3}ze^{i\alpha} - 1$, $\varphi_2(z) = \frac{1}{3}ze^{i\alpha} + 1$, where $\alpha = r\pi$ with irrational r . Let us show that convex invariant set $\tilde{K}$ with respect to the system φ_1, φ_2 has infinite set of sides. As it follows from the Proposition 1, for any compact set $A \subset \mathbb{R}^2$, $\lim_{n \rightarrow \infty} \tilde{T}^n(A) = \tilde{K}$.

Let $P_0 = \{\bar{0}\}$ and $P_n = \tilde{T}(P_0)$. Obviously, each P_n is a convex polygon and $P_0 \subset P_1 \subset \dots \subset P_n \subset \dots$, so $\tilde{K} = \lim_{n \rightarrow +\infty} P_n = \bigcup P_n$.

It's easy to see that n -th polygon P_n has n pairs of opposite sides l'_k, l''_k , each having length $2/3^{k-1}$ and forming an angle $(k-1)\alpha$ with horizontal axis:

It's obvious for $n = 2$. Assuming the fact is true for P_n , observe that $P_{n+1} = P_n + l_{n+1}$, where l_{n+1} is a line segment $\left[-\frac{e^{ik\alpha}}{3^k}, \frac{e^{ik\alpha}}{3^n} \right]$ and $A + B$




Figure 1:

denotes the set $\{a + b, a \in A, b \in B\}$. Since neither of sides l'_k, l''_k of P_n is parallel to l_{n+1} , P_{n+1} has two more sides equal to l_{n+1} . The sides $l'_k(P_n)$, $l''_k(P_n)$ are equal and parallel for different n and converge to the sides $l'_k(\tilde{K})$ and $l''_k(\tilde{K})$ of $\tilde{K}$ having the same length and direction. So $\tilde{K}$ has the sides l'_k and l''_k for each $k \in N$. Taking their sum we see that the length of $\partial\tilde{K}$ is more or equal to $4 + \frac{4}{3} + \dots + \frac{4}{3^n} + \dots = 6$.




Figure 2:

To get the opposite inequality, consider the sets $D_n = \tilde{T}^n(D)$, where $D = \{x^2 + y^2 \leq 1\}$ is an unit disc of radius $\rho > 3$. They form a nested sequence $D_1 \supset D_2 \supset \dots \supset D_n \supset \dots$ and boundary ∂D_n of D_n consists of n pairs of sides l'_k, l''_k , each having length $2/3^{k-1}$ and forming an angle $(k-1)\alpha$ with horizontal axis and of $2n$ arcs of radius $\rho/3^{n+1}$, so it has total length $\frac{2\pi\rho}{3^{n+1}} + 4 + \frac{4}{3} + \dots + \frac{4}{3^n}$. As n tends to ∞ we obtain the opposite inequality $H^1(\partial\tilde{K}) \leq 6$, so $H^1(\partial\tilde{K}) = 6$. In other words, 1-dimensional measure of $\partial\tilde{K}$ equals the sum of lengths of sides of $\tilde{K}$. Therefore the set of vertices of $\tilde{K}$ has 1-dimensional Hausdorff measure 0.

3 Open convex set condition.

DEFINITION 2 . A system $\{\varphi_1, \dots, \varphi_n\}$ satisfies open convex set condition (OCSC), if there is non-empty open convex set O , such that

- (i) $\cup \varphi_i(O) \subset O$,
- (ii) $\varphi_i(O) \cap \varphi_j(O) = \emptyset$ if $i \neq j$.

PROPOSITION 3 . If the interior U of the set $\tilde{K}$ is non-empty, then

- (i) $\cup \varphi_i(U) \subset O$,
- (ii) $\varphi_i(U) \cap \varphi_j(U) = \emptyset$ if $i \neq j$.

Proof. The first inclusion is obvious. Let now O be the open set from the OCS condition. Since $\tilde{T}(\bar{O}) \subset \bar{O}$, the set $\tilde{K}$ is contained in $\bar{O}$. Therefore $\varphi_i(\tilde{K}) \cap \varphi_j(\tilde{K}) \subset \varphi_i(\bar{O}) \cap \varphi_j(\bar{O}) \subset \varphi_i(\bar{O} \setminus O) \cup \varphi_j(\bar{O} \setminus O)$. The latter set is closed and nowhere dense in $\mathbb{R}^2$ and the set $\varphi_i(U) \cap \varphi_j(U)$ is it's open subset. Therefore it is empty.

LEMMA 4 . Let $U_1, U_2, \dots, U_n$ be closed convex domains with disjoint interiors and let V be the convex hull of the set $\bigcup_{i=1}^n U_k$. Let $Q_i = U_i \cap \partial V$, $Q_{i,s}$ be the components of the set Q_i , and m_i be the number of such components. Then $\sum_{i=1}^n m_i \leq 2n - 2$.

Proof. The case $n = 2$ is obvious, because $m_1 = m_2 = 1$.




Figure 3:

Suppose it is already proved for any family consisting of $n - 1 \geq 2$ sets U_i . If all $m_i = 1$, the statement also holds, so we can suppose that for some Q_i , the number of it's components $m_i \geq 2$. Denote the components of $V \setminus U_i$ by N_k , $k = 1, \dots, m_i$. Each of the sets U_j , $j \neq i$ is contained in the closure of some component N_k , and each N_k contains at least one of the sets U_j , $j \neq i$: therefore $m_i \leq n - 1$.

Let n_k be the number of all $U_j \subset \bar{N}_k$. Obviously, $1 \leq n_k \leq n - 2$.

Consider a family $\{U_i\} \cup \{U_j | U_j \subset \bar{N}_k\}$ and its convex hull $\bar{N}_k \cup U_i$. The number of the sets U_k in this family is $n_k + 1 \leq n - 1$, so we can apply the statement of Lemma and obtain $1 + \sum_{U_j \subset \bar{N}_k} m_j \leq 2n_k$. A total sum over all components N_k gives the desired inequality

$$m = \sum_{j \neq i} m_j + m_i \leq 2 \sum n_k = 2(n - 1).$$

PROPOSITION 5 . 1) Each of the sets $Q_i, i = 1 \dots, n$ contains finite number of components $Q_{i,s}$, and their total number is less or equal to $2n - 2$.
 2) If $n > 2$ then for $i \neq j$ the set $Q_{i,s} \cap Q_{j,t}$ is either empty or is a point .
 3) If a sequence of indices $(i_1, \dots, i_n)$ is an initial interval of $(j_1, \dots, j_m)$, then $Q_{i_1, \dots, i_n} \subset Q_{j_1, \dots, j_m}$. If neither of sequences $(i_1, \dots, i_n), (j_1, \dots, j_m)$ is an initial interval of the other then $Q_{i_1, \dots, i_n} \cap Q_{j_1, \dots, j_m}$ is either empty or is a point.
 4) If for some j , $Q_j = \emptyset$, then for each sequence $(i_1, \dots, i_n)$, $Q_{i_1, \dots, i_n, j} = \emptyset$

Proof. The first statement results from the previous lemma.

To prove the second statement, suppose the set $Q_{i,s} \cap Q_{j,t}$ contains two points x, y . Let l be a line segment with endpoints x, y . If $l \subset Q_{i,s} \cap Q_{j,t}$, then some half-neighbourhood of the point $\frac{x+y}{2}$ is contained in $\tilde{K}_i \cap \tilde{K}_j$, which contradicts OCSC. By the very reason $l \setminus \{x, y\}$ cannot be contained neither in $\tilde{K}_i$ nor in $\tilde{K}_j$. Since $n > 2$, l cannot be contained in the intersection of the boundaries of the sets $\tilde{K}_i$ and $\tilde{K}_j$.

The third statement follows directly from 2). 4) is obvious.

4 Finiteness of the set of sides of $\tilde{K}$.

DEFINITION 6 . We call a side l with endpoints $x_1, x_2 \in F$ a side of order 1, if there are such $j_1 \neq j_2$ that $x_1 \in Q_{j_1}, x_2 \in Q_{j_2}$

PROPOSITION 7 . If $\{\varphi_1, \dots, \varphi_n\}$ satisfies OCSC, then the set of sides of order 1 is finite.

Proof. The sides of order 1 are the closures of components of $\partial \tilde{K} \setminus \bigcup_{i,s} Q_{i,s}$ so from Proposition 5 it follows that there is no more than $2n - 2$ such sides.

PROPOSITION 8 . Each side $l' \subset \partial \tilde{K}$ is an image $\varphi_{i_1 i_2 \dots i_p}(l)$ of some side l of order 1.

Proof Let x', y' be the endpoints of the side l' . If l' is contained in a component $Q_{i_1 i_2 \dots i_p}$, then $\varphi_{i_1 i_2 \dots i_p}^{-1}(l') \subset \partial \tilde{K}$. It follows from the statement 3) of the Proposition 5, that among all components $Q_{i_1 i_2 \dots i_p}$, containing the side l' , there is a component $Q_{i_1 i_2 \dots i_p}$ of maximal order. There exist different $j, k = 1, \dots, n$, such that $x' \in Q_{i_1 i_2 \dots i_p j}$, $y' \in Q_{i_1 i_2 \dots i_p k}$. Then the side $l = \varphi_{i_1 i_2 \dots i_p}^{-1}(l')$ is an side of order 1, because it's endpoints $x = \varphi_{i_1 i_2 \dots i_p}^{-1}(x')$ and $y = \varphi_{i_1 i_2 \dots i_p}^{-1}(y')$ belong to different components Q_j and Q_k .

THEOREM 9 *If $\varphi_1, \dots, \varphi_n$ satisfies OCSC and all the angles $\alpha_1, \dots, \alpha_n$ are commensurable with π , then the set of sides of $\tilde{K}$ is finite.*

Proof. Let θ be the greatest common divisor of the angles $\alpha_1, \dots, \alpha_n, \pi$. Let $l_1, \dots, l_m$ be the edges of order 1, and $\beta_1, \dots, \beta_m$ be the angles formed by $l_1, \dots, l_m$ and the axis Ox . Each of the sides of $\tilde{K}$ is an image $\varphi_{i_1 \dots i_p}(l_j)$ of some side l_j of order 1, and therefore the angle it forms with the horizontal axis is of the form $\beta_j + n\theta$. The set of all such angles is finite, so the set of all sides of $\tilde{K}$ is finite. ■

4.1 Vertices and corner points.

Since every vertex z_0 of $\tilde{K}$ lies in $\partial \tilde{K} \cap K$ there is an infinite sequence $K_{i_1} \supset K_{i_1 i_2} \supset \dots K_{i_1 i_2 \dots i_p} \supset \dots$ such that $z_0 = \bigcap_{p=1}^{\infty} K_{i_1 i_2 \dots i_p}$. So for each p there is a vertex $z_p = \varphi_{i_1 \dots i_p}^{-1}(z_0)$. Thus each vertex z_0 has an infinite sequence of predecessors $z_p \in \partial \tilde{K}$ such that $z_0 = \varphi_{i_1 \dots i_p}(z_p)$ for some $\varphi_{i_1 \dots i_p}$.

The vertex is *periodic* if there is a periodic sequence $i_1, i_2, \dots, i_p, \dots$ defining z_0 . In this case one of z_0 's predecessors, say z_m , is a fixed point of some $\varphi_{i_1 \dots i_p}$. We call a vertex z_0 a *corner point* if right and left tangents to $\tilde{K}$ at z_0 do not coincide.

PROPOSITION 10 . *Each corner point $z_0 \in \partial \tilde{K}$ is a periodic vertex and both right and left tangents at a corner point z_0 are sides l_0^+ and l_0^- .*

Proof. Let θ_0 be an angle between right and left tangents to $\tilde{K}$ at z_0 , and θ_p an angle between right and left tangents to $\tilde{K}$ at z_p . Since $\varphi_{i_1 \dots i_p}(z_p) = z_0$, the angle θ_p is more or equal to θ_0 . The sum of θ_p over all of different predecessors of z_0 does not exceed 2π , therefore the sequence $z_1, z_2, \dots, z_p, \dots$ contains no more than $2\pi/\theta_0$ different elements. This shows that z_0 is periodic.

Let now $V^+(z_0)$ denote a half-neighborhood of the point z_0 in $\partial \tilde{K}$ taken in positive direction. As we see from Proposition 5 for each p there is unique p -tuple $i_0, i_1, \dots, i_p$ for which $Q_{i_0 i_1 \dots i_p} \cup V^+(z_0) \setminus z_0$ is non-empty for each $V^+(z_0)$. The sequence $Q_{i_0} \supset Q_{i_0 i_1} \supset \dots Q_{i_0 i_1 \dots i_p} \supset \dots$ defines unique sequence $z_1, z_2, \dots, z_p, \dots$ of predecessors of z_0 having the additional property that

for each half-neighborhood $V^+(z_0)$ of z_0 and $V^+(z_p)$ of z_p the intersection $V^+(z_0) \setminus z_0 \cap \varphi_{i_0 i_1 \dots i_p}(V^+(z_p))$ is non-empty. Since the number of different z_p 's is finite, one of them, say z_m , is a fixed point of some $h = \varphi_{j_1 \dots j_q}$, satisfying $h(V^+(z_p)) \subset V^+(z_p)$. The latter is possible only when $V^+(z_p)$ is a straight line interval.

PROPOSITION 11 . *If all the angles $\alpha_1, \dots, \alpha_n$ are such that for each set of non-negative integers $k_1, \dots, k_n$, $k_1 \alpha_1 + \dots + k_n \alpha_n \neq m\pi, m \in \mathbb{N}$, then the set F of the vertices of $\tilde{K}$ is infinite.*

Proof. If the set of vertices of $\tilde{K}$ is finite, all of them are corner points. Since none of $\varphi_{i_1 \dots i_q}$ has rotation angle $m\pi$, $\partial\tilde{K}$ has no corner points, so the set F is infinite.

4.2 The main theorem.

We call two systems and $(\varphi'_1, \dots, \varphi'_n)$ *convex equivalent*, if they generate the same convex invariant set $\tilde{K}$.

Let $S' = (\varphi'_1, \dots, \varphi'_n)$ be p -th refinement of the system $S = (\varphi_1, \dots, \varphi_n)$, i.e. the set of all mappings $\varphi_{i_1 \dots i_p}$ of order p . The system S' is convex equivalent to S , and S' satisfies OCSC if S does.

PROPOSITION 12 . *For any system of similitudes $(\varphi_1, \dots, \varphi_n)$ satisfying the OCS condition, there is a convex equivalent system $(\psi_1, \dots, \psi_n)$ satisfying the OCS condition such that*

$$\text{all the sets } \psi_{i_1 \dots i_k}(\tilde{K} \cap \partial\tilde{K}) \text{ are connected} \quad (1)$$

$$\forall i \quad \varphi_i(\tilde{K}) \cap F = \emptyset \quad (2)$$

$$\forall i \neq j \quad \varphi_i(\tilde{K}) \cap F \not\subset \varphi_j(\tilde{K}) \cap F. \quad (3)$$

Proof. Suppose for some i the set $Q_i = \tilde{K}_i \cap \partial\tilde{K}$, is not connected, and $Q_{i,1} \dots Q_{i,s}$ are it's components. Let δ_i be minimal distance between different components of Q_i . Let δ be the smallest of all δ_i among non-connected Q_i 's.

With each component $Q_{i,k}$ of Q_i we associate a set $N_{i,k}$ of all unit outer normal vectors to $\partial\tilde{K}$ at points $x \in Q_{i,k}$. If any edge $l \subset \partial\tilde{K}$ has both of it's edges in Q_i then it lies completely in one of it's components, say $Q_{i,k}$.

Therefore for each i , the sets $N_{i,k}$ are disjoint closed arcs on the unit circle. Let θ_i be the length of the shortest complementary arc to $\bigcup_k N_{i,k}$. Let θ be the smallest of all θ_i 's.

Choose such p_0 that $q^{p_0} \cdot \text{diam}(\tilde{K}) < \delta$.

For any p_0 -tuple $i_1 \dots i_{p_0}$ the set $\varphi_{i_1 \dots i_{p_0}}(\tilde{K})$ has diameter less than δ , so the set $\varphi_{i_1 \dots i_{p_0}}(\tilde{K})$ may have non-empty intersection with at most one of the components of Q_{i_1} . Therefore $Q_{i_1 \dots i_{p_0}}$ lies completely in some component of Q_{i_1} . Take any sequence $(j_1 \dots j_q)$. By the same reason, for any $k \leq q$ the set $Q_{j_1 \dots j_p i_1 \dots i_{p_0}}$ must be contained completely in some component of $Q_{j_1 \dots j_k}$. Consider $S' = (\varphi'_1, \dots, \varphi'_{n'})$ be p -th refinement of the system $S = (\varphi_1, \dots, \varphi_n)$. By the above argument each non-empty set $Q_{ij} = \psi_{ij}(\tilde{K}) \cup \partial\tilde{K}$ is contained in a unique component of the set $Q'_{ij} = \psi_i(\tilde{K}) \cup \partial\tilde{K}$. Replacing S by S' if necessary, we may suppose from that moment that for each i, j , Q_{ij} is contained completely in some component of Q_i .

Assume some $Q_{i_1 \dots i_p}$ is non-connected and $Q_1^{(1)}$ and $Q_2^{(1)}$ are it's two adjoining components, joined by an arc $\Delta^{(1)}$ in $\partial\tilde{K}$ with endpoints $\xi^{(1)} \in Q_1^{(1)}$ and $\eta^{(1)} \in Q_2^{(1)}$. Then for each $1 \leq k \leq p$ the set $Q_{i_k \dots i_p}$ is also non-connected and contains two components $Q_1^{(k)}, Q_2^{(k)}$ satisfying $\varphi_{i_1 \dots i_{k-1}}(Q_i^{(k)}) \cap \partial\tilde{K} = Q_i^{(1)}$ also joined in $\partial\tilde{K}$ by an arc $\Delta^{(k)}$ so that $\varphi_{i_1 \dots i_{k-1}}(\Delta^{(k)}) \cap \partial\tilde{K} = \Delta^{(1)}$.

Let $N(\Delta^{(k)})$ be the set of all unit outer normal vectors to $\partial\tilde{K}$ at points $x \in \Delta^{(k)}$. All these sets are open arcs. Since φ_i preserve angles between vectors, these arcs have the same length. They cannot coincide for $k_1 \neq k_2$ because the diameters of corresponding sets $\Delta^{(k_1)}$ and $\Delta^{(k_2)}$ are different. They cannot have non-empty intersection, because of the OCSC condition. Therefore there cannot be more than $\frac{2\pi}{\theta}$ of such arcs.

So, if $p_1 > \frac{2\pi}{\theta}$, then each of the sets $Q_{i_1 \dots i_{p_1}}$ is connected.

Therefore the p_1 -th refinement of the system S satisfies conditions 1) and OCSC.

A system $(\varphi_i, \dots, \varphi_n)$ of contraction maps satisfying conditions (1), (2), (3), and OCSC, will be called *regular*.

LEMMA 13 *If the system $(\varphi_i, \dots, \varphi_n)$ is regular, then the number of all components $Q_{i_1 \dots i_p} \subset \partial\tilde{K}$ of order p , having non-empty interior in $\partial\tilde{K}$ does not exceed n^p .*

Proof. The statement is obvious for $p = 1$.

Suppose it's true for all components $Q_{i_1 \dots i_{p-1}}$ of order $p - 1$.

Suppose a component $Q_{i_1 \dots i_{p-1}}$ contains a p -component $Q_{i_1 \dots i_{p-1}j}$ different from $Q_{i_1 \dots i_{p-1}}$. Then the endpoints ξ_j, η_j of the component Q_j must satisfy either $\varphi_{i_1 \dots i_{p-1}}(\xi_j) \in \overset{\circ}{Q}_{i_1 \dots i_{p-1}}$ or $\varphi_{i_1 \dots i_{p-1}}(\eta_j) \in \overset{\circ}{Q}_{i_1 \dots i_{p-1}}$.

Using this observation we can estimate the number of all components of order p $Q_{i_1 \dots i_p}$ having non-empty interior in $\partial\tilde{K}$.

Consider the sets of unit normal vectors $N_{i_1 \dots i_p}^\circ = \bigcup_{x \in \overset{\circ}{Q}_{i_1 \dots i_p}} N_x$, where

$\overset{\circ}{Q}_{i_1 \dots i_p}$ is the interior of the component $Q_{i_1 \dots i_p}$ in $\partial\tilde{K}$. If the set $\overset{\circ}{Q}_{i_1 \dots i_p}$ is

non-empty, then $N_{i_1 \dots i_p}^\circ$ is an open arc of the unit circle. It is clear that $N_{i_1}^\circ \supset N_{i_1 i_2}^\circ \supset \dots \supset N_{i_1 \dots i_p}^\circ \supset \dots$ and $N_{i_1 \dots i_p}^\circ = \varphi_{i_1 \dots i_p}(N_{i_p}^\circ) \cap N_{i_1 \dots i_{p-1}}^\circ$.

If the set $N_{i_1 \dots i_p}^\circ$ differs from $N_{i_1 \dots i_{p-1}}^\circ$ one of the endpoints of the arc $N_{i_1 \dots i_p}^\circ$ must lie in $N_{i_1 \dots i_{p-1}}^\circ$.

So, if $\beta_{i_p}^-$ and $\beta_{i_p}^+$ are the endpoints of the arc $N_{i_p}^\circ$, then one of the following inequalities hold:

$$\beta_{i_1 \dots i_p}^- < \alpha_{i_1} + \dots + \alpha_{i_{p-1}} + \beta_{i_p}^- < \beta_{i_1 \dots i_p}^+$$

$$\beta_{i_1 \dots i_p}^- < \alpha_{i_1} + \dots + \alpha_{i_{p-1}} + \beta_{i_p}^+ < \beta_{i_1 \dots i_p}^+$$

The sum $\alpha_{i_1} + \dots + \alpha_{i_{p-1}}$ is the same for different permutations of $i_1 \dots i_{p-1}$, whereas the sets $N_{i_1 \dots i_{p-1}}^\circ$ are disjoint. Therefore we can take the union of all those $N_{i_1 \dots i_{p-1}}^\circ$ for which $\alpha_{i_1} + \dots + \alpha_{i_{p-1}} = \gamma$, and denote it by N_γ° . The union of the interiors $\overset{\circ}{Q}_{i_1 \dots i_{p-1}}$ of the corresponding $p-1$ -components in $\partial \tilde{K}$ we denote by Q_γ° . Let p_γ be the number of these components.

Thus, if a component $\overset{\circ}{Q}_{i_1 \dots i_p} \subset P_\gamma$ is different from $\overset{\circ}{Q}_{i_1 \dots i_{p-1}}$, then one of the conditions $\gamma + \beta_{i_p}^- \in N_\gamma^\circ$, $\gamma + \beta_{i_p}^+ \in N_\gamma^\circ$ must hold.

More exactly, if $p-1$ -component $\overset{\circ}{Q}_{i_1 \dots i_{p-1}}$ contains $q+1$ p -components $Q_{i_1 \dots i_p}, \dots, Q_{i_1 \dots i_{p-1}(i_p+q)}$, then the system of inequalities

$$\begin{cases} \beta_{i_1 \dots i_{p-1}}^- < \alpha_{i_1} + \dots + \alpha_{i_{p-1}} + \beta_{i_p+k}^+ < \beta_{i_1 \dots i_{p-1}}^+ \\ \beta_{i_1 \dots i_{p-1}}^- < \alpha_{i_1} + \dots + \alpha_{i_{p-1}} + \beta_{(i_p+k+1)}^- < \beta_{i_1 \dots i_{p-1}}^+ \end{cases}$$

holds for $k = 0, \dots, q-1$.

Therefore if m is a total number of those angles β_j^+ and β_j^- , for which the conditions $\gamma + \beta_j^- \in N_\gamma^\circ$ or $\gamma + \beta_j^+ \in N_\gamma^\circ$ hold, then the number of components of order p contained in Q_γ is not greater than $p_\gamma + m$.

The number of different sets Q_γ , is in its turn no greater than the number of different summands in the expansion of $(x_1 + \dots + x_n)^{p-1}$, which is equal to $\frac{(n+p-2)!}{(p-1)!(n-1)!}$. Therefore the number of different components (having in $\partial \tilde{K}$ non-empty interior) of order p exceeds the number of order $(p-1)$ components by a number not greater than $n \cdot \frac{(n+p-2)!}{(p-1)!(n-1)!}$.

So the total number of components of order p is less or equal to

$$n \left(1 + \frac{n!}{1!(n-1)!} + \frac{(n+1)!}{2!(n-1)!} + \dots + \frac{(n+p-2)!}{(p-1)!(n-1)!} \right) = \frac{(n+p-1)!}{(p-1)!(n-1)!},$$

which is less than p^{n+1} .

THEOREM 14 *If the system $(\varphi_1 \dots \varphi_n)$ satisfies the OCS condition, then Hausdorff dimension of the set F of the vertices of $\partial \tilde{K}$ is zero.*

Proof. We can suppose the system $(\varphi_1 \dots \varphi_n)$ is regular. The union of all $Q_{i_1 \dots i_p}$ consisting of one point is at most countable, so we may consider the set $F' = F \setminus \cup Q_{i_1 \dots i_p}$, which has the same Hausdorff dimension as F .

For each p , the set F' is covered by components $Q_{i_1 \dots i_p}$, having non-empty interior in $\partial \tilde{K}$. The total number of such components does not exceed p^{n+1} , and the diameter of each of them is less or equal to $q^p \cdot \text{diam}(\tilde{K})$. Since $\lim_{p \rightarrow \infty} \left(-\frac{\ln p^{n+1}}{\ln q^p} \right) = 0$, Hausdorff dimension of the set F' , and, therefore of F , is zero [1].

COROLLARY 15 *If the system $\varphi_1, \dots, \varphi_n$ satisfies OCSC and all the angles $\alpha_1, \dots, \alpha_n$ are commensurable with π , then the set $\tilde{K}$ is a convex finite polygon.*

Proof. Since the number of sides of $\tilde{K}$ (Theorem 9) is finite, the set F has finite number of components. All they have zero measure therefore each of them is a point.

References

- [1] *Falconer K.J.:* Fractal geometry:mathematical foundations and applications . – J.Wiley and Sons, New York, 1990.
- [2] *Hutchinson J.:* Fractals and self-similarity. – Indiana Univ. Math. J., 30, No 5, 1981, pp.713-747.